

Theoretical Physics Institute
University of Minnesota

TPI-MINN-92/67-T
UND-HEP-92-BIG 07
NSF-ITP-92-156
December 1992

A QCD “Manifesto” on Inclusive Decays of Beauty and Charm¹

I.I.Bigi

Dept. of Physics, University of Notre Dame du Lac, Notre Dame, IN 46556

B.Blok

Institute for Theoretical Physics, University of California at Santa Barbara, Santa Barbara, CA 93106

M.Shifman

Theoretical Physics Institute, University of Minnesota, Minneapolis, MN 55455

N.G.Uraltsev

Dept. of Physics, University of Notre Dame du Lac, Notre Dame, IN 46556

and

St.Petersburg Nuclear Physics Institute, Gatchina, St.Petersburg 188350, Russia²

A.Vainshtein

Theoretical Physics Institute, University of Minnesota, Minneapolis, MN 55455

and

Budker Institute for Nuclear Physics, Novosibirsk 630090, Russia

Abstract

A selfconsistent treatment of nonperturbative effects in inclusive nonleptonic and semileptonic decays of beauty and charm hadrons is presented. It is illustrated by calculating semileptonic branching ratios, lifetime ratios, radiative decay rates and the lepton spectra in semileptonic decays.

¹talk presented at DPF meeting of APS, November 1992

²permanent address

1. Sketch Of General Method

It is the goal of our analysis to calculate transition rates for inclusive non-leptonic and semileptonic decays in terms of the fundamental parameters, namely quark masses, KM parameters and Λ_{QCD} without invoking Voodoo. Nonperturbative effects – namely boundstate dynamics in the initial state and hadronization in the final state – have to be incorporated here. We will do that via an expansion in $1/m_Q$ (m_Q is the mass of the heavy flavour quark Q). For the beauty quark mass is certainly much larger than typical hadronic scales, and an extrapolation down to charm hopefully still makes sense.

The width for the decay of a hadron with the heavy flavor Q into an inclusive final state X is obtained from the transition operator $\hat{T}(Q \rightarrow X \rightarrow Q)$ describing the forward scattering of a heavy quark Q via an intermediate state X . Then we take the imaginary part of the matrix element of \hat{T} between the state H_Q :

$$\Gamma(H_Q \rightarrow X) \propto G_F^2 \langle H_Q | \text{Im} \hat{T}(Q \rightarrow X \rightarrow Q) | H_Q \rangle. \quad (1)$$

One actually proceeds in two steps to evaluate this quantity:

(a) The nonlocal second-order operator \hat{T} is expanded into a series of local operators $O^{(i)}$ of increasing dimension with coefficients c_i that contain increasing powers of $1/m_Q$

$$\text{Im} \hat{T}(Q \rightarrow X \rightarrow Q) \simeq \sum_i c_i O^{(i)} \quad (2)$$

(Strictly speaking, the expansion parameter is $1/E$ where E is the energy release, rather than $1/m_Q$.) The lowest dimensional operators $O^{(i)}$ thus dominate for large m_Q .

(b) One has to evaluate the matrix elements $\langle H_Q | O^{(i)} | H_Q \rangle$; it is at this level that one differentiates between the decays of heavy flavor baryons and neutral and charged pseudoscalar mesons, i.e. Λ_Q , P^0 and P^\pm . This approach was explicitly formulated in ref.¹ and has been supported by the analysis of ref.².

(1) The general expansion of $\text{Im} \hat{T}$ is as follows (for details see ¹⁻⁴)

$$\text{Im} \hat{T}(Q \rightarrow X \rightarrow Q) = \tilde{c}_3^{(X)} m_Q^5 \bar{Q}Q + \tilde{c}_5^{(X)} m_Q^3 \bar{Q} i\sigma G Q + \tilde{c}_6^{(X)} m_Q^2 (\bar{Q}\Gamma q)(\bar{q}\Gamma Q) + \mathcal{O}(m_Q) \quad (3)$$

with $i\sigma G = i\gamma_\mu \gamma_\nu G_{\mu\nu}$, $G_{\mu\nu}$ being the gluonic field strength tensor. The coefficients $\tilde{c}_i^{(X)}$ are calculable dimensionless numbers that depend on the mode X . Once the matrix elements between the states H_Q are formed the first term on the right hand side of eq.(3) yields (among other things) the usual free ‘spectator’ result; the second one constitutes a new type of ‘spectator’ contribution; the third term represents ‘non-spectator’ contributions due to W-exchange (‘WA’) and interference effects (‘PI’).

(2) The relevant matrix elements of the operators $\bar{Q}Q$, $\bar{Q} i\sigma G Q$ and $(\bar{Q}\Gamma q)(\bar{q}\Gamma Q)$ have been evaluated so far. For $\bar{Q}Q$ one employs a nonrelativistic $1/m_Q$ expansion:

$$\bar{Q}Q = v_\mu \bar{Q} \gamma_\mu Q + \frac{1}{4m_Q^2} \bar{Q} i\sigma G Q - \frac{1}{2m_Q^2} \bar{Q} (D^2 - (v \cdot D)^2) Q + \mathcal{O}(1/m_Q^3) \quad (4)$$

D_μ denotes here the covariant derivative and v_μ is the 4-velocity vector. One finds:

$$\langle H_Q | \bar{Q} \gamma_\mu Q | H_Q \rangle = 1 \cdot v_\mu \quad \left(\text{if } \langle H_Q(\vec{p}') | H_Q(\vec{p}) \rangle = \frac{E}{M} \cdot (2\pi)^3 \delta^3(\vec{p}' - \vec{p}) \right) \quad (5)$$

$$\langle P_Q | \bar{Q} i\sigma G Q | P_Q \rangle \simeq \frac{3}{2} (M^2(V_Q) - M^2(P_Q)) \quad (6)$$

$$\langle \Lambda_Q | \bar{Q} i\sigma G Q | \Lambda_Q \rangle \simeq 0; \quad \Lambda_Q = \Lambda_b, \Lambda_c; \quad P_Q = B, D; \quad V_Q = B^*, D^*. \quad (7)$$

The operator $\bar{Q}(D^2 - (vD)^2)Q$ describes the kinetic energy of the heavy quark in the gluon background field; the masses of Λ_Q , P_Q , V_Q yield some information on its size.

Finally we employ the approximation of factorization $\langle P_Q | (\bar{Q}\Gamma q)(\bar{q}\Gamma Q) | P_Q \rangle \simeq \langle P_Q | \bar{Q}\Gamma q | 0 \rangle \langle 0 | \bar{q}\Gamma Q | P_Q \rangle$ with the operators normalized at the low hadronic scale.

2. Qualitative Results

The decay widths of a heavy flavor hadron H_Q look as follows:

$$\Gamma(H_Q) = \Gamma(Q) \cdot [1 + a_2^{(H_Q)}/m_Q^2 + a_3^{(H_Q)}/m_Q^3 + \mathcal{O}(1/m_Q^4)] \quad (8)$$

where $\Gamma(Q)$ is the free quark width with all purely perturbative effects included.

- (a) The free spectator ansatz thus holds through terms of order $1/m_Q$.
- (b) The first nonperturbative corrections enter on the $1/m_Q^2$ level with

$$a_2^{(\Lambda_b)} \neq a_2^{(B^-)} = a_2^{(B_d)} \simeq a_2^{(B_s)} \quad (9)$$

- (c) ‘Conventional’ non-spectator effects – WA and PI – emerge on the $1/m_Q^3$ level:

$$a_3^{(B^-)} < a_3^{(B_d)} < a_3^{(\Lambda_b)} \quad (10)$$

- (d) In charm decays these nonperturbative corrections exhibit similar pattern while being larger by roughly factors of $(m_b/m_c)^2 \sim 10$ and $(m_b/m_c)^3 \sim 30$, respectively.

3. (Semi-)Quantitative Phenomenology

3.1 Semileptonic Branching Ratios: Corrections $1/m_b^2$

The semileptonic and nonleptonic transition operators for $b \rightarrow c$ decays in the external gluon field are given by:

$$\hat{\Gamma}_{SL} = F \left\{ z_0 \bar{b} b - \frac{z_1}{m_b^2} \bar{b} i\sigma G b \right\}, \quad \hat{\Gamma}_{NL} = FN_C \left\{ A_0 (z_0 \bar{b} b - \frac{z_1}{m_b^2} \bar{b} i\sigma G b) - \frac{4A_2 z_2}{m_b^2} \bar{b} i\sigma G b \right\} \quad (11)$$

with $z_0 = 1 - 8x + 8x^3 - x^4 - 12x^2 \log(x)$, $z_1 = (1-x)^4$, $z_2 = (1-x)^3$ and $x = m_c^2/m_b^2$ describing the relative phase spaces, $A_0 = (c_+^2 + c_-^2)/2 + (c_+^2 - c_-^2)/2N_C$ and $A_2 = (c_+^2 - c_-^2)/2N_C$ denoting the QCD radiative corrections (N_C =number of

colours) and $F = G_F^2 m_b^5 |V_{cb}|^2 / 192\pi^3$. Thus we find a reduction in the semileptonic branching ratio:

$$\frac{\delta BR_{SL}(B)}{BR_{SL}(B)} \simeq 6 \cdot \frac{M_{B^*}^2 - M_B^2}{m_b^2} \frac{A_2 z_2}{A_0 z_0} \cdot BR_{NL}(B) < 0 \quad (12)$$

This reduction is of the order of 5% in B decays; $BR_{SL}(\Lambda_b)$ remains practically unaffected by nonperturbative corrections on the $1/m_Q^2$ level due to eq.(7).

In charm decays on the other hand one obtains a much larger reduction:

$$\frac{\delta BR_{SL}(D)}{BR_{SL}(D)} \sim -\mathcal{O}(50\%) \quad . \quad (13)$$

It is worth noting that in charmed particles some of the $\mathcal{O}(m_c^{-3})$ corrections are numerically as large as the $\mathcal{O}(m_c^{-2})$ ones.

The last term in eq.(11) is produced by an interference between two different color flow amplitudes (it represents a non-factorizable contribution). It is worth noting that it is approximately saturated⁴ by non-factorizable contributions to a few exclusive amplitudes that were estimated in ref.⁵ arising due to the same operator.

3.2 Lifetime Ratios

Small differences are predicted for the lifetimes of the different beauty hadrons:

$$1.02 \leq \tau(B^-)/\tau(B_d) \leq 1.08 \quad \text{for} \quad 130\text{MeV} < f_B < 250\text{MeV} \quad ; \quad (14)$$

these effects are mainly due to ‘PI’ with ‘WA’ being rather small^{1,2}. A preliminary analysis shows that $\tau(\Lambda_b)$ and $\tau(B)$ could differ by up to 10-15 %.

Extrapolating these findings down to charm one obtains much bigger effects: $\tau(D^+)/\tau(D^0) \sim 2$ mainly due to ‘PI’ with ‘WA’ providing a 20% contribution at most. An intriguing ‘roller coaster’ story thus unfolds behind the semileptonic branching ratios for D mesons: on the m_Q^5 and m_Q^4 levels one gets the parton result $BR_{SL}(D) \simeq 15\%$; $1/m_Q^2$ corrections suppress it roughly by a factor of two: $BR_{SL}(D) \simeq 8\%$; $1/m_Q^3$ effects finally differentiate between D^0 and D^+ decays by suppressing $\Gamma_{NL}(D^+)$ such that $BR_{SL}(D^+) \simeq 16\%$ holds while affecting $BR_{SL}(D^0)$ only a little.

3.3 $b \rightarrow s\gamma$

The chromomagnetic dipole operator affects also radiative $b \rightarrow s + \gamma$ decays:

$$\hat{\Gamma}_\gamma \propto m_b^5 \{ \bar{b}b - 1/m_b^2 \bar{b}i\sigma G b \} \quad ; \quad (15)$$

yet the ratio $BR(B \rightarrow \gamma + (S = -1)) / BR(B \rightarrow l\nu X)$ remains practically unchanged.

3.4 Lepton Spectra in Semileptonic Decays

Lepton spectra in semileptonic beauty and charm decays can be treated⁶ in an analogous fashion where one relies on an expansion in $1/(p_Q - p_l)^2$ rather than

in $1/m_Q^2$ with p_Q [p_l] denoting the momentum of the heavy quark Q [the lepton l]. In the limit of the vanishing quark mass in the final state of $b \rightarrow ql\nu$ and neglecting the gluon bremsstrahlung one obtains⁷:

$$\frac{d\Gamma}{dy} \propto y^2 \left[\frac{3}{2} - y + \left(\frac{5}{3}y + \frac{1}{3}\delta(1-y) + \frac{1}{6}(2y^2 - y^3) \delta'(1-y) \right) \frac{K}{2} + \left(2 + \frac{5}{3}y - \frac{11}{6}\delta(1-y) \right) \frac{G}{2} \right] \quad (16)$$

where $y = 2E_l/m_b$, $K = 1/m_b^2 \cdot \langle H_b | \bar{b} \vec{D}^2 b | H_b \rangle$ and $G = 1/2m_b^2 \cdot \langle H_b | \bar{b} i\sigma G b | H_b \rangle$. The δ -functions (for $m_q = 0$) reflect the blowup of the expansion near the endpoint. It is important however that the integral over a small region $\Delta y > \mu/m_b$ (μ is a hadronic scale) around the endpoint is correctly reproduced by that of eq.(16).

4. Outlook

Obviously our analysis on the lepton spectra has to be finalized⁷. Beyond that there are two intriguing areas for further study, namely

- the question of $SU(3)_{flavour}$ breaking, e.g. the relation between D^0 and D_s and between B_d and B_s decays;
- the vast realm of heavy flavour baryon decays.

5. Acknowledgements

A.V. is grateful for helpful discussions to L.McLerran and M.Voloshin. This work was supported in part by the NSF under grant number NSF-PHY 92 13313 and in part by DOE under grant number DOE-ER-40105-1100.

6. References

1. M.Voloshin and M.Shifman, Sov.J.Nucl.Phys. **41** (1985) 120; Sov.Phys.-JETP **64** (1986) 698.
2. I.I.Bigi and N.G.Uraltsev, *Phys.Lett.* **B280** (1992) 271.
3. I.I.Bigi, N.G.Uraltsev and A.Vainshtein, *Phys.Lett.* **B293** (1992) 430.
4. B.Blok and M.Shifman, preprints NSF-ITP-92-103, -115.
5. B.Blok and M.Shifman, preprint NSF-ITP-92-76.
6. The first detailed discussion of semileptonic spectra was given in: J.Chay, H.Georgi, B.Grinstein, *Phys.Lett.* **B247** (1990) 399.
7. I.I.Bigi, M.Shifman, N.G.Uraltsev and A.Vainshtein, in preparation.